Appl. No.

09/804,457

Filed

March 12, 2001

AMENDMENTS TO THE CLAIMS

1. (Presently Amended): A method of characterizing the biological activity of a candidate compound comprising:

exposing one or more cells to said compound;

repetitively exposing said one or more cells to a series of two or more electric fields so as to effect a change in transmembrane potential of said one or more cells without using a patch clamp, wherein said transmembrane potential changes predominantly in a single direction away from a starting transmembrane potential over the course of said series of electric fields <u>due to a continuing and additive accumulation of charge in said cell over the course of said series of electric fields</u>; and

monitoring, without using a patch clamp, changes in the transmembrane potential of said one or more cells to characterize the biological activity of said compound.

- 2. (Original) The method of Claim 1, wherein said monitoring comprises detecting fluorescence emission from an area of observation containing said one or more cells.
 - 3. (Original) The method of Claim 1, wherein said electric fields are biphasic.
- 4. (Original) The method of Claim 3, additionally comprising limiting spatial variation in electric field intensity so as to minimize irreversible cell electroporation.
- 5. (Original) The method of Claim 1, wherein one or more electrical fields cause an ion channel of interest to cycle between different voltage dependent states.
- 6. (Original) The method of Claim 5, wherein said one or more electrical fields cause an ion channel of interest to open.
- 7. (Original) The method of Claim 5, wherein said one or more electrical fields cause an ion channel of interest to be released from inactivation.
- 8. (Original) The method of Claim 1, wherein said one or more cells comprise a voltage sensor selected from the group consisting of a FRET based voltage sensor, an electrochromic transmembrane potential dye, a transmembrane potential redistribution dye, an ion sensitive fluorescent or luminescent molecule and a radioactive ion.

Appl. No. : 09/804,457 Filed : March 12, 2001

9. (Original) The method of Claim 1, wherein said one or more cells comprise a voltage regulated ion channel.

- 10. (Original) The method of Claim 9, wherein said voltage regulated ion channel is selected from the group consisting of a potassium channel, a calcium channel, a chloride channel and a sodium channel.
- 11. (Original) The method of Claim 1, wherein said electric field exhibits limited spatial variation in intensity in the area of observation of less than about 25% from a mean intensity in that area.
- 12. (Original) The method of Claim 11, wherein said one or more electrical fields varies over an area of observation by no more than about 15 % from the mean electrical field at any one time.
- 13. (Original) The method of Claim 12, wherein said one or more electrical fields varies over an area of observation by no more than about 5 % from the mean electrical field at any one time.
- 14. (Original) The method of Claim 1, wherein said one or more electrical fields comprises stimulation with either a square wave-form, a sinusoidal wave-form or a saw tooth wave-form.
- 15. (Original) The method of Claim 1, wherein said one or more electrical fields have an amplitude within the range of about 10 V/cm to about 100 V/cm.
- 16. (Original) The method of Claim 15, wherein said one or more electrical fields have an amplitude within the range of about 20 V/cm to about 80 V/cm.
- 17. (Original) The method of Claim 1, wherein said one or more electrical fields are repeated at a frequency of stimulation that is greater than or equal to the reciprocal of the transmembrane time constant of said one or more cells.
- 18. (Original) The method of Claim 1, wherein said one or more electrical fields are repeated at a frequency of stimulation within the range of zero to 1kHz.
- 19. (Original) The method of Claim 1, wherein said one or more electrical fields have a pulse duration within the range of about 100 microseconds to about 20 milliseconds.
- 20. (Original) The method of Claim 1, wherein said transmembrane potential is developed across the plasma membrane of said one or more cells.

Appl. No.

09/804,457

Filed

March 12, 2001

21. (Presently Amended) A method of assaying the biochemical activity of a compound against a target ion channel comprising:

selecting a cell line having a normal resting transmembrane potential corresponding to a selected voltage dependent state of said target ion channel;

expressing said target ion channel in a population of cells of said selected cell line;

exposing said population of cells to said compound;

repetitively exposing said population of cells to a series of two or more electric fields so as to effect a change in transmembrane potential of said population of cells, wherein said transmembrane potential changes predominantly in one direction away from a starting transmembrane potential over the course of said series of electric fields <u>due to a continuing and additive accumulation of charge in said cell over the course of said series of electric fields</u>; and

monitoring changes in the transmembrane potential of said population of cells to characterize the biochemical activity of said compound.

- 22. (Previously Amended) The method of Claim 21, wherein said target ion channel is endogenously expressed in the cell line.
- 23. (Original) The method of Claim 21, wherein said cell line is transfected with nucleic acid encoding said target ion channel.
- 24. (Previously Presented) The method of Claim 23, wherein said cell line expresses substantially only said target ion channel.
- 25. (Original) The method of Claim 24, wherein said cell line is selected from the group consisting of CHL, LTK(-), and CHO-K1.
- 26. (Original) The method of Claim 21 wherein said target ion channel is a sodium channel, and wherein said population of cells is selected from the group consisting of CHL cells, LTK(-) cells, and CHO-K1 cells.
- 27. (Original) The method of Claim 21 wherein said target ion channel is a sodium channel, and wherein said population of cells is selected from the group consisting of HEK-293 cells, RBL cells, F11 cells, and HL5 cells.

Appl. No. : 09/804,457 Filed : March 12, 2001

28. (Original) The method of Claim 21 wherein said target ion channel is a potassium channel, and wherein said population of cells is selected from the group consisting of CHL cells, LTK(-) cells, and CHO-K1 cells.

29. (Original) The method of Claim 21 wherein said target ion channel is a calcium channel, and wherein said population of cells is selected from the group consisting of CHL cells, LTK(-) cells, and CHO-K1 cells.

30-48 Cancelled

- 49. (Previously Presented) The method of Claim 1, additionally comprising using said electric fields to maintain said transmembrane potential within a predefined range.
 - 50. (Previously Presented) The method of Claim 21, additionally comprising using said electric fields to maintain said transmembrane potential within a predefined range.

Appl. No. Filed

09/804,457

March 12, 2001

Summary of Interview

The applicant thanks the Examiner for the time taken during the telephone interview. The differences between the continuing voltage change produced by the series of electric fields of the invention and electric field induced action potentials was discussed.